

Peatlands and the Historic Environment

An Introduction to their Cultural and Heritage Value



Summary

This document introduces the archaeological, palaeoenvironmental and cultural significance and value of peatlands. It describes where peat deposits are found and the wealth and diversity of heritage assets associated with them – illustrated with examples from both upland and lowland peat settings. Peatlands have further public value as carbon stores and for their rich biodiversity; their importance straddles the natural environment and historic environment sectors. This document outlines the main natural and human-induced threats facing peatlands, of common concern to both sectors, and summarises some of the ways in which peatlands can be protected.

This document has been prepared by Zoë Hazell and Jim Williams (Historic England) with contributions from Benjamin Gearey (University College Cork) and Deborah Land (Natural England).

This edition published by Historic England September 2024.

All images © Historic England unless otherwise stated.

Please refer to this document as:

Historic England 2024 *Peatlands and the Historic Environment: An Introduction to their Cultural and Heritage Value.* HEAG300a (version 2) Swindon. Historic England.

HistoricEngland.org.uk/advice/technical-advice/peatlands/

Front cover: The Holme Fen post, Holme Fen National Nature Reserve (NNR), Cambridgeshire. In 1850 the post was put into place with its pointed top at ground level, to show and monitor the peat loss resulting from draining nearby Whittlesey Mere (see Hutchinson 1980; Waltham 2000). This original post (together with a more-recent adjacent one) is recognised for its historic environment value, being Grade II Listed (List Entry Number 1288011). September 2020 © Zoë Hazell

Contents

1.	Introduction	1
2.	The resource	3
	2.1 Peat resource2.2 Historic environment resource	
3.	The cultural and heritage value	7
	 3.1 The deposits 3.2 Historic environment features 3.3 Historic landscapes 3.4 Wider cultural heritage 	12 16
4.	Managing peatlands	18
	4.1 Threats to peatlands4.2 Peat restoration works4.3 'Doing nothing'	20
5.	Designations and protections	21
6.	Further reading	22
7.	References	23
8.	Glossary	26
9.	Where to get advice	29
10.	Acknowledgements	30

1. Introduction

Background and context

Peatlands are at risk of loss from a range of threats, both direct and indirect. With the increased awareness of their natural and cultural-heritage capitals and values, works are being undertaken across the country to protect and restore peatlands that have been impacted by degradation and erosion. Some of the primary drivers for restoration relate to climate change – carbon storage, water regulation and ecology/biodiversity – but re-establishing healthy ecosystem function of peatlands is also necessary to benefit the historic environment. Peatlands contain a wealth of heritage assets, archaeological remains and palaeoenvironmental information, which – through ensuring peatland longevity – will also be preserved.

In 2021 Defra published the England Peat Action Plan (EPAP) (UK Government 2021), setting out plans for restoring England's peatlands and facilitated by a central government funding scheme. Crucially, the plan highlighted the need for close working across natural and historic environment organisations. This was to ensure that peatland restoration delivers maximum public value; it is essential that the significance and contribution that historic environment elements of peatlands make to their overall value, is recognised. In direct response to the England Peat Action Plan, this document was produced, together with:

- Peatland restoration and the historic environment: Standards for delivering environmentally sustainable peatland restoration projects (Natural England and Historic England 2022), and
- *Peatlands and the Historic Environment: Guidance for Carrying out Peat Restoration* HEAG300b (Historic England, forthcoming)

Definitions

The term 'historic environment' includes the concepts of: historic landscapes (as created, modified and managed by human actions), individual heritage features on the surface, archaeology (above, within and below the peat), records of past environmental changes (as preserved in/as the peat), cultural identity and living heritage. The term 'peatland' is used here to refer to both/either upland and/or lowland deposits.

The definitions of peats follow the classifications of Natural England (2010, 7): i) deep peaty soils at >40cm deep, ii) shallow peaty soils at 10–40cm deep, and iii) soils with peaty pockets.

Other specific term definitions are included in a Glossary at the end of the document.

2. The resource

Peatlands preserve unique archaeological and palaeoecological records, are living historic landscapes, and are part of our biocultural heritage. Peat deposits are important archives of past human activities and environments, often forming over thousands of years. The waterlogged conditions that characterise peatlands result in the exceptional preservation of natural and cultural organic remains, unrivalled at typical 'dryland' archaeological sites.

Peat deposits form through the gradual, upward accumulation of organic matter. This buildup of plant matter occurs because waterlogged, anoxic conditions prevent the remains from decomposing. Peat initiation and development are influenced by a site's physical and climatic setting (e.g. topography, water input) and by human activities. Evidence of anthropogenic influences on peatlands extends back to prehistoric times.

Crucially, many areas of peat deposits no longer look like the bogs or fens they once were, particularly where they have been altered by extensive drainage, peat extraction or agricultural regimes.

2.1 Peat resource

It is estimated that peatlands total an area of around 3 million hectares (12.2% of total UK land area), of which around 640,000ha (22%) is thought to be in a near-natural condition (Evans et al 2017). In England, the area of mapped peat is estimated to be around 682,230ha, however, 186,000ha of this is mapped as wasted peat (Natural England 2010) (Figure 1).

There are three main formation-types of peatland in the UK:

- blanket bog in the uplands and far north;
- rarer lowland raised bogs; and
- fen peatlands.




Figure 1: Map of the UK showing the distribution of different types of peatlands (after Natural England 2010, 9, map 2).

Peatlands can be small, a single site or feature, or large encompassing whole landscapes. Some of the oldest peatlands in England date back to the early Holocene period c. 12,000 years ago. In temperate regions peat depths of more than 10m are not uncommon, with accumulation rates equivalent to 1cm per 10 years (for example, Walton Moss, Cumbria (Figure 2)). Because of the way they are formed peatlands are important for capturing and storing carbon. Therefore, looking after them is important for tackling the climate crisis.

Peatland classifications, typologies and nomenclatures are complex; for the purpose of this document, the term peatland is used to refer to both/either upland and/or lowland peat deposits.



Figure 2: The lowland raised bog at Walton Moss, Cumbria is (unlike its near neighbour Bolton Moss) a near pristine raised bog. It is widely regarded as the most intact ombrotrophic bog in England (Natural England 2015a, 16). It is a SSSI, with areas also designated as NNR and SAC. As such, it has been the subject of multiple palaeoenvironmental studies (e.g. Hughes et al 2000). Image taken December 2013 © Deborah Land.

2.2 Historic environment resource

Historic environment features are present at/on the surface, within, and below the peat. Quantifying the extent and distribution of peatland archaeology is particularly challenging. This is because much of the resource is buried, and as yet unknown. Where figures are known, these are for areas which have experienced greatest disturbance (particularly through peat cutting) and therefore most concentrated recording. Whilst the figures also represent the most damage/loss of archaeological remains, these are still likely to be underestimates given the amounts of peat removed/lost prior to archaeological recording.

Results of regional archaeological surveys in England have been used to suggest (minimum) estimates of: 2.2 sites per km² in lowland peatlands (i.e. 4,200 sites) and at least 1.2 sites per km² in upland peatlands (i.e. 1,800 sites) (Van de Noort et al 2002). But there are also regional differences; extrapolating data from the Brue Valley and Sedgemoor in the Somerset Levels, would put the estimated number of sites in lowland peatlands higher, at more than 3.4 sites per km² (i.e. 7,000).

These figures are based on the results of multiple wetland surveys funded by Historic England. As well as the Somerset Levels, areas studied extensively include: the East Anglian Fens, the Humber wetlands and North West England. Projects exploring the condition and management of England's wetlands as a whole include the Monuments at Risk in England's Wetlands (MAREW) and Heritage Management of England's Wetlands (HMEW) projects.

3. The cultural and heritage value

There are multiple facets to the cultural and heritage value of peatlands. They are important archives of archaeological and palaeoenvironmental records, preserved within – and as – the peat deposits themselves. Peatlands were part and parcel of life. They are historic landscapes and part of cultural identities, with an intangible heritage associated with their past use and exploitation. For more information see Gearey et al (2011), Gearey and Chapman (2023) and Flint and Jennings (2020).

3.1 The deposits

Organic remains can be preserved in peatlands for thousands of years, due to the waterlogged, anoxic conditions that characterise these deposits. Peatlands are important archives of our past, documenting climate and environmental change, and providing evidence of human activities that took place in, and influenced, these alterations (see Gearey et al 2011; Charman 2002). These data are important for understanding contemporary processes (such as climate change) and for providing the ecological 'baseline' context of peatland restoration processes.

Archaeological and palaeoenvironmental remains are frequently discussed and considered in terms of their research 'potential'.

Archaeology

Direct evidence of human occupation and activity (archaeological remains) can be found:

- at/on the peat surface,
- within the peat, and
- below the peat.

Examples include structures (such as wooden trackways (Figure 3), platforms (Figure 4) and dwellings), domestic and personal items (Figure 5a and 5b), tools and implements, ceremonial monuments (e.g. stone rows (Figure 6a and 6b), burial chambers (Figure 7a and 7b) and cairns (Figure 8a)), rock art (Figure 8b), food (e.g. the Celtic/fava (*Vicia faba*) beans at Glastonbury Lake Village (Reid 1917)), and of the people themselves (1.e. 'bog bodies') (see Stevens and Chapman 2020).

Figure 3 (top):

Excavating the Neolithic timber Lindholme trackway, Hatfield Moor, South Yorkshire, 2004–2006. Note the more-recent drainage ditch cutting across the trackway itself © Henry Chapman.

Figure 4 (bottom):

The central platform (the largest of the lake edge platforms) under excavation in 2013, at the Early Mesolithic site of Star Carr, North Yorkshire © Star Carr Project, CC BY-NC 4.0 (see Milner et al 2018a, b).





Figure 5: (a) Star Carr has yielded the largest assemblage of Mesolithic bone and antler artefacts in Europe, with artefact types unique to the site. Here, the *in situ* red deer frontlet (possible headdress/hunting disguise) with a smaller roe deer crania (see Milner et al 2018a). © Star Carr Project.

(b) A partially charred birch bark
roll from Star Carr; one of a total of
161 excavated from the site over the
years. Possible uses include as tapers/
torches, in tar production, and/or as
fishing floats (see Milner et al 2018b).
© Paul Shields, University of York.







Figure 6: Examples of surface and within-peat archaeology: the Neolithic stone row at Cut Hill, Dartmoor, Devon (Fyfe and Greeves 2010); (a) showing the line of stones visible at the surface, and (b) and the submerged, horizontal stone [with the short scale bar placed on top], together with the peat profile sampled for palaeoecological analyses and radiocarbon dating. © Ralph Fyfe.

Figure 7: The Bronze Age cist (burial chamber), Whitehorse Hill, Dartmoor, Devon (see Jones 2017). Within the cist were preserved the remains of a cremation burial. Finds included: beads, a worked flint, textiles, basketry, and wooden studs, all wrapped up within a bear pelt. *Filipendula* sp. (meadowsweet) pollen was identified from samples within the cist, suggesting that a floral tribute was included as part of the burial rite; (a) the tin, amber, shale and ceramic beads, and the wooden *Euonymus* sp. (spindle wood) (possible ear) studs, and (b) excavating the cist and sampling the adjacent peat profile. Both images © Dartmoor National Park Authority.







Figure 8: Archaeological remains exposed after the 2003 fire on Fylingdales moor, North Yorkshire. Note that in January 2006, when this photograph was taken, the vegetation was recovering. (a) Three probably prehistoric features: a cup-marked stone in the foreground, a small stone cairn just behind (likely to be an Early Bronze Age burial monument), and what is probably a clearance cairn in the background; (b) an example of the prehistoric rock art (probably late Neolithic). © Jonathan Last, Historic England.

Palaeoecology

The peat deposits themselves are made up of remains of the plants and animals that once lived in and on the peat, and can be used to reconstruct a site's environmental history and provide an understanding of the context in which human activity occurred. The palaeoecological data can also be used to understand regional and global climate and environmental change.

As well as providing direct information on past vegetation (through the analysis of e.g. leaves, seeds, pollen and spores) peat also contains proxy indicators of past site conditions (e.g. testate amoebae and insects) and evidence of past burning events (e.g. micro-charcoal) (see Historic England's Environmental Archaeology guidance). Where such deposits remain intact, they provide continuous records of past environmental histories. The faster the accumulation rates, the higher the resolution of the environmental record.

Scientific dating

The main dating techniques used on peat deposits and organic archaeological remains are radiocarbon dating and dendrochronology. The technique that uses the lead isotope ²¹⁰Pb can be used to date more-recent deposits (<200 years old), and, where available, tephra layers can be used as chronological markers. In addition, archaeological artefacts found within features and associated with activities, can provide a good indication of when these events took place and their duration.

Other analyses

Analysis of other isotopes (e.g. oxygen, hydrogen, carbon and nitrogen) can provide information on past environmental conditions, particularly precipitation and wetness. Evidence of past industrial activities and pollution over a range of timescales can be determined from SCPs (Spheroidal Carbonaceous Particles) and atmospheric metal deposition records.

3.2 Historic environment features

Within these landscapes sit historic environment features (heritage assets). Whilst many are directly related to the use/exploitation of the peat itself, not all are. As well as being important in their own right, their significance and value derive from their wider, contextual (i.e. landscape) setting as outlined above.

Individual features are associated with:

- historical peat working (extraction): the peat cuttings themselves (domestic, industrial) which can vary in size and extent (Figure 9a and 9b) e.g. the landscape-scale Norfolk Broads; baulks, banks and ditches, peat stacks, steadings (Figure 10); tramways (Figure 11); hollow-way routes to and between cuttings; associated buildings e.g. for processing and storing peat (Figures 12 and 13) e.g. for petrochemicals such as naphtha;
- past water management regimes e.g. grips/ditches, and wind pumps (Figure 14);
- exploitation of other resources e.g. reed/sedge beds for thatching, and wildfowl and eels for food;
- occupation e.g. houses, field systems;
- industrial archaeology e.g. tin, coal, lead mining tinners' buildings, extraction shafts, spoil heaps, leats, associated buildings (Figure 15);
- standing buildings, associated with other activities e.g. transport infrastructure; and
- military activities e.g. 20th century built structures, trenches, wartime aeroplane crash sites (Figure 16a and 16b).

Many aspects of peatland landscapes are still not fully understood or recorded, with upland features in particular being discovered as walkover surveys are undertaken prior to peat restoration works.



Figure 9: (a) Peat cuttings (likely post-medieval) in the foreground, on the upland hills at Exe Plain, Exmoor, Somerset, in February 2012 (by D. Grady) © Historic England Archive (ref. 27413_002). (b) Chapman's Broad (centre), adjacent to the larger Hickling Broad, Norfolk, in March 1938. They are part of The Norfolk Broads, created as the result of extensive Medieval peat cutting; as demonstrated by Dr Joyce Lambert in the 1950s (e.g. Lambert and Smith 1960; also see Natural England 2015b, Moss 2001, 11). © Historic England Archive (ref. EPW056549).



Figure 10: Remains of a steading (rick) on Bodmin Moor, Cornwall – a low platform within a shallow ditch on which cut turves were dried. This historic asset was only identified in advance of peatland restoration works, despite being in an area that has previously undergone extensive archaeological research and survey. This feature type is thought to be specific to this region. © Southwest Peatland Partnership.



Figure 11: Remains of a tramway at Amicombe, Dartmoor surveyed as part of peat restoration works funded by Defra and South West Water. Sleepers can still be found in place within the vegetation. © Southwest Peatland Partnership.



Figure 12: A ruined and part-demolished peat charring oven at Rattlebrook Head Peat Works, Dartmoor, installed in the late 1870s. © Phil Newman.





Figure 13: Grade II listed remains of the late 18th to early 19th century Old Gang Peat Store, North Yorkshire. (List Entry Number 1131503). The peat was used as fuel to smelt the locally-mined lead. © Historic England Archive (ref. 28813_046) taken September 2015 (D. MacLeod)

Figure 14:

View across Sedge Fen (Wicken Fen, Cambridgeshire) showing the historic, Grade II listed (List Entry Number 1161096) (re-located) wind-pump, and the visitor centre. The windpump dates from the early 20th century and was originally used to drain water for peat cutting on Adventurer's Fen, but was moved in 1956. February 2014 © Zoë Hazell.



Figure 15: Evidence of coal mining and associated processing on Fountains Fell, North Yorkshire, part of which is a Scheduled Monument (ref. 1017715). © Historic England Archive (ref. 34119_001) taken September 2019 (E. Trevarthen)





Figure 16: Remains of the WWII Spitfire X4593 from Holme Fen, Cambridgeshire which crashed on 22 November 1940 during a training flight. The body of Pilot Officer Harold Penketh was recovered at the time, and the remains of the aeroplane were excavated in 2015 by Oxford Archaeology East. The photographs show the propeller *in situ* (a), and excavated (b). All photos © Oxford Archaeology.

3.3 Historic landscapes

On a holistic, landscape scale, peatland landscapes ('peatscapes') provide composite evidence of being shaped, modified and/or created by past human activities. These extend from the influence of prehistoric human activities on early peatland development, to more recent effects of large-scale historic drainage/extraction schemes.

3.4 Wider cultural heritage

Peatlands also have links to intangible heritage and cultural identity, for example:

- the history and heritage of peat cutting, for example the tools used and livelihoods supported, as presented for example at Nidderdale Museum, North Yorkshire (https:// www.nidderdalemuseum.com), and the Kirbuster Museum, Orkney (https://www. orkney.com/listings/kirbuster-museum);
- the traditional language associated with peat deposits (e.g. Finlay MacLeod's "Some Lewis Moorland Terms: A Peat Glossary", in Macfarlane 2016); and oral histories (e.g. the "Moorlands: People, Places, Stories" project in Upper Nidderdale, North Yorkshire (see Nidderdale AONB 2018));
- associative value for example, Wicken Fen, Cambridgeshire was where ecologists and botanists Sir Harry Godwin, Sir Arthur Tansley and Dr Max Walters, as well as Charles Darwin, undertook their pioneering research (see Hazell 2014);
- as a resource-base for past uses, that might not have left lasting evidence in the landscape, for example harvesting *Sphagnum* moss for medical dressings;
- the landscapes as represented in art (e.g. Thomas Wade's Turf Cutters (1869); John William Inchbold's Peat Burning (c 1864–6); Joseph Beuys; Fred Ingrams) and in literature (e.g. Seamus Heaney; Edna O'Brien); and
- as a continued resource today e.g. in whisky manufacture (see Godsman and Van De Greindt 2008).

Key points

Peatlands preserve unique, evocative and fragile archaeological sites and artefacts that are generally not found on drylands, and provide an amazing record of environmental and landscape change. They make it possible to place people's activities within their environmental setting. Their extensive cultural and heritage connections provide exceptional opportunities for public enjoyment and engagement.

4. Managing peatlands

Good management of peatlands is vital in order to protect and maintain their multiple public benefits. Increasingly, this is being driven by the urgent need to stem growth in atmospheric carbon dioxide concentrations to avoid catastrophic climate change. Peatlands are a crucial store of carbon – reducing peat loss, and maintaining and restoring healthy peatlands, are essential for global carbon reduction and can also help reduce the impacts of an already changing climate. Overall, concerns for the loss of peatlands are shared by the natural environment and historic environment sectors. Therefore, managing peatland sites requires close collaboration and joint working.

4.1 Threats to peatlands

Peatlands as a whole are at risk of loss from a range of direct and indirect threats. Peat loss occurs on a range of scales and rates, from incremental losses to drastic events, and their impact is cumulative, putting heritage above, within and below the peat at risk (see Cover image). The main factors driving peat loss include: climate change, agriculture, tree planting, root damage, fires (including managed burning), peat extraction, dewatering, desiccation, erosion, and land drainage (see Gearey et al 2011 for more detail). Many of these are interlinked, and will increase in frequency and/or severity as a result of climate change, for example, the incidence of fires. Some threats are more relevant for certain peatland types; for example, arable agriculture is a particular threat to lowland peats. Holman (2009, 9, 14) used various sources to estimate that deep peats within the East Anglian Fens were wasting at rates of up to 2.1cm per year (for land described as 'intensive arable'). Further to these, development threats (such as windfarm construction) and their associated infrastructure (e.g. access roads) can alter hydrological function and directly affect peat deposits. Changes in land use, such as to paludiculture, can also affect peat deposits, with variability between crops and their root/rhizome depths.

Whilst peat loss is unwanted and needs to be reversed and/or reduced, the destructive processes involved can lead to important archaeological discoveries (Figures 4, 7a, 7b, 8a and 8b) and provide opportunities for engagement (Figure 17).



Figure 17: Peatlands offer incredible opportunities for communities to explore and engage with the archaeological and palaeoecological resource. The lowland peatland of Hatfield Moors has been the focus of concerted public engagement via The Heritage Lottery Council; funding The Isle of Axholme and Hatfield Chase Landscape Partnership (IOAHC) (https://ioahc.net/). Events have been held on the themes of: (a) experimental archaeology – the project Trackway Through Time built a reconstruction trackway, including making the stone tools and felling the tree, and (b) palaeoecology – Project Wildscape's 'Palaeofest' event included coring to retrieve samples, followed by a pollen citizen science workshop. © Benjamin Gearey.



As well as the threats to peatlands (as outlined above), further specific threats to the historic environment include:

- changes to site hydrogeology; for example, fluctuating water tables, and water-table draw-down resulting from drainage (which can cause degradation to archaeological and palaeoenvironmental remains, e.g. see Davies et al 2015); and
- actions that can change and/or obscure surface features, and alter their wider setting including works undertaken to restore peat areas.

4.2 Peat restoration works

In order to stabilise and protect peat deposits at risk of loss (e.g. through erosion) peat restoration works are undertaken. However, restoration works themselves can result in changes to, and loss of, historic environment features, archaeological remains and peat deposits if not carefully planned. Methods used in restoration schemes include: reprofiling, grip blocking (using dams of peat, wood, plastic sheeting, or heather/coir plant bales to block up old drainage ditches) and vegetation planting/seeding (e.g. of *Sphagnum* moss). The method(s) used need to be tailored depending on the character of each site, and will need to consider loss of, or disturbance to, the sediments, as well as peat compression. More information is provided in Peatlands and the Historic Environment: Guidance for Carrying out Peat Restoration (forthcoming).

4.3 'Doing nothing'

In some settings 'doing nothing' is a direct threat to the preservation of peatlands' historic environment features and deposits, resulting in permanent loss of the finite historic environment resource. Remedial actions (i.e. restoration works) are therefore necessary, and enable an initial limited, managed and recorded loss; in preference to the alternative which would be a continued, uncontrolled and unrecorded loss.

Peatlands also play an important role in the climate system through carbon storage and emissions; a recent Greenhouse Gas (GHG) emission estimate of UK peatlands is around 23 million tons carbon dioxide equivalent per year (Evans et al 2017). Restoring the natural functioning of peatlands will help reduce this.

Key points

Overall, concerns for the loss of peatlands are shared by the natural and historic environment sectors. Peatlands are at risk of loss from a range of direct and indirect threats, and increasingly, pro-active management approaches are required to protect them. Together, these can be viewed as opportunities to discover and record historic environment features as they are revealed.

Designations and protections

Sites and features are recognised and protected through historic environment designations (e.g. scheduling and listing) and/or natural environment designations (e.g. SSSI, NNR, LNR and SAC). Yet designations do not necessarily protect sites from loss, especially those in settings that are drying out. Some specific site types will have additional statutory protections e.g. aircraft crash sites. Sites that sit within protected landscapes (e.g. National Parks, and National Landscapes) are afforded protection as part of the overarching management responsibilities of the governing authority. Other land management schemes (e.g. environmental stewardship) also help ensure that land-owners manage their natural and cultural assets appropriately.

International designations are also relevant, for example, World Heritage Site status (administrated by UNESCO), and the Ramsar Convention that designates wetlands of international importance.

It is essential to recognise that natural and historic environment designations have different management priorities and approaches. These do not always directly align, and highlight the importance of close cross-sector working and engagement.

6. Further reading

The Historic England-funded wetland surveys are all published; The Somerset Levels Project, the Fenland Project, the Humber Wetlands Project, and the North West Wetlands Survey.

Guidance on specific aspects of archaeology and heritage are available through the Historic England website, in particular:

Environmental Archaeology https://historicengland.org.uk/advice/technical-advice/archaeologicalscience/ environmental-archaeology/

More on peatlands, their form and function can be found in Natural England publications. For example, see England's peatlands: Carbon storage and greenhouse gases NE257, available at: http://publications.naturalengland.org.uk/publication/30021

The International Union for Conservation of Nature (IUCN) UK Peatland Programme website also hosts a range of peat resources, including the comprehensive historic environment review by Gearey et al (2011); see: https://www.iucn-uk-peatlandprogramme.org/

7. References

Charman, D 2002 Peatlands and Environmental Change Chichester. John Wiley and Sons, Ltd

Davies, H, Fyfe, R and Charman, D 2015 'Does peatland drainage damage the palaeoecological record?' *Review of Palaeobotany and Palynology* 221, 92–105

English Heritage 2008 Conservation Principles, Policies and Guidance: For the Sustainable Management of the Historic Environment https://historicengland.org.uk/images-books/ publications/conservation-principles-sustainable-management-historic-environment/

Evans, C, Artz, R, Moxley, J, Smyth, M-A, Taylor, E, Archer, N, Burden, A, Williamson, J, Donnelly, D, Thomson, A, Buys, G, Malcolm, H, Wilson, D, Renou-Wilson, F and Potts, J 2017 *Implementation of an emissions inventory for UK peatlands* Report to the Department for Business, Energy and Industrial Strategy. Bangor. Centre for Ecology and Hydrology https:// uk-air.defra.gov.uk/library/reports?report_id=980

Flint, A and Jennings, B 2020 'Saturated with meaning: peatlands, heritage and folklore' *Time and Mind* 13(3), 283–305

Fyfe, R and Greeves, T 2010 'The date and context of a stone row: Cut Hill, Dartmoor, southwest England' *Antiquity* 84, 55–70

Gearey, B, Bermingham, N, Chapman, H, Charman, D, Fletcher, W, Fyfe, R, Quartermaine, J and Van de Noort, R 2011 *Peatlands and the Historic Environment* Scientific review. Final version, June 2011. Produced for the IUCN UK Peatland Programme

Gearey, B and Chapman, H 2023 *An Introduction to Peatland Archaeology and Palaeoenvironments* Studying Scientific Archaeology 6. Oxford. Oxbow Books

Godsman, NM and Van De Griendt, NA 2008 'The importance of peat for whisky production', in *After Wise Use – The Future of Peatlands* Proceedings of the 13th International Peat Congress: Peat In Energy, 106–108

Hazell, Z 2014 *Wicken Fen: Statement of Significance* Draft report for Historic England Project 6240 Historic England forthcoming *Peatlands and the Historic Environment: Guidance for Carrying out Peat Restoration* HEAG300b Swindon: Historic England

Holman, IP 2009 *An estimate of peat reserves and loss in the East Anglian Fens* Unpublished report for the RSPB by the Department of Natural Resources, Cranfield University

Hughes, PDM, Mauquoy, D, Barber, KE and Langdon, PG 2000 'Mire development pathways and palaeoclimatic records from a full Holocene peat archive at Walton Moss, Cumbria, England' *The Holocene* 10(4), 465–479

Hutchinson, JN 1980 'The record of peat wastage in the East Anglian fenlands at Holme Post, 1848–1978AD' *Journal of Ecology* 68, 229–249

Jones, AM (ed.) 2017 Preserved in the Peat Oxbow Books

Lambert, JM and Smith, CT 1960 'The Norfolk Broads as man-made features' *New Scientist* 7(176), 775–777, 31 March

Macfarlane, R 2016 Landmarks Penguin Books

Milner, N, Conneller, C and Taylor B (ed.s) 2018a *Star Carr. Volume 1: a persistent place in a changing world* White Rose University Press https://doi.org/10.22599/book1

Milner, N, Conneller, C and Taylor B (ed.s) 2018b *Star Carr. Volume 2: studies in technology, subsistence and environment* White Rose University Press https://doi.org/10.22599/book2

Moss, B 2001 The Broads [New Naturalist 89] London. HarperCollins

Natural England 2010 *England's peatlands: Carbon storage and greenhouse gases* NE257 http://publications.naturalengland.org.uk/publication/30021

Natural England 2015a *Bolton Fell and Walton Mosses SSSI Cumbria*. Notification under section 28C of the Wildlife and Countryside Act 1981. Supporting Information

Natural England 2015b *National Character Area Profile: 80 The Broads* NE449, April 2015 http://publications.naturalengland.org.uk/publication/11549064

Natural England and Historic England 2022 *Peatland restoration and the historic environment: Standards for delivering environmentally sustainable peatland restoration projects* https://historicengland.org.uk/images-books/publications/peatland-restorationhistoric-environment/ Nidderdale AONB 2018 *Moorlands: People Places Stories*. Upper Nidderdale Landscape Partnership http://uppernidderdale.org.uk/wp-content/uploads/2018/11/Moorlands_ booklet_WEB.pdf

Reid, C 1917 'Plants, wild and cultivated', pp. 625–630 in Bulleid, A and Gray, HSG (ed.s) *The Glastonbury Lake Village. A Full Description of the Excavations and the Relics Discovered*, 1892–1907, Vol. II. Glastonbury. Glastonbury Antiquarian Society

Stevens, J and Chapman, H 2020 'Regional patterns in bog body distribution – a case study from the UK' *Journal of Wetland Archaeology* 19(1–2) [JWA Bog Bodies 2019 Special Edition], 131–153

UK Government 2021 *England Peat Action Plan* May 2021 https://assets.publishing.service. gov.uk/government/uploads/system/uploads/attachment_data/file/1010786/england-peataction-plan.pdf

Van de Noort, R, Fletcher, W, Thomas, G, Carstairs, I and Patrick, D 2002 *Monuments at Risk in England's Wetlands* Unpublished report by The University of Exeter, on behalf of English Heritage, version 3

Walker, M, Johnsen, S, Olander Rasmussen, S, Popp, T, Steffensen, J-P, Gibbard, P, Hoek, W, Lowe, J, Andrews, J, Björck, S, Cwynar, LC, Hughen, K, Kershaw, P, Kromer, B, Litt, T, Lowe, DJ, Nakagawa, T, Newnham, R and Schwander, J 2009 'Formal definition and dating of the GSSP (Global Stratotype Section and Point) for the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records' *Journal of Quaternary Science* 24(1), 3–17

Waltham, AC 2000 'Peat subsidence at the Holme Post' Mercian Geologist 15(1), 49–51

8. Glossary

Anoxic - oxygen-depleted

Blanket bog – an expanse of peat covering extensive areas of upland hills, whose water input is primarily from rainfall ('ombrotrophic'). It is an overarching term used to describe a complex of feature sub-types.

Dendrochronology – tree ring dating. This is based on the annual growth patterns of most tree species in temperate regions

Desiccation – drying out

Dewatering - the intentional removal/loss of ground/surface water

Fen – a peatland which receives water input from groundwater and/or runoff ('minerotrophic') (Charman 2002, 6)

Greenhouse Gases (GHGs) – peatlands release a selection of GHGs; CO_2 – carbon dioxide, CH_4 – methane, and N₂O – nitrous oxide

Grips - artificial ditch features dug to drain peat

Holocene – the current warm period following the end of the last glacial c 12,000 years ago (Walker et al 2009)

Isotopes – are atoms of an atomic element that have a different number of neutrons (but the same number of protons and electrons) e.g. ${}^{13}C/{}^{12}C$, ${}^{18}O/{}^{16}O$

LNR – Local Nature Reserve

Micro-charcoal – microscopic charcoal fragments produced during burning events. They are used as evidence of past fires

Naphtha – a flammable liquid produced from the distillation of peat

National Landscapes – previously known as AONBs (Areas of Outstanding Natural Beauty) and still legally designated as such (see https://national-landscapes.org.uk/)

NNR – National Nature Reserve

NPPF – National Planning Policy Framework; the "government's planning policies for England and how these are expected to be applied"; see https://www.gov.uk/government/ publications/national-planning-policy-framework--2

Palaeoecological – palaeoecology = 'old ecology'; palaeoecological refers to the study of past environments from the material evidence surviving of those past ecosystems; plants, animals, sediments etc

Paludiculture – farming of commercial crops in rewetted peatlands

²¹⁰Pb dating – a radiometric dating technique that uses this radioactive isotope of lead

Peat [classifications] – in England, peat deposits are classified by Natural England (2010, 7) as:

- Deep peaty soils: areas covered with a majority of peat >40cm deep
- Shallow peaty soils: areas with a majority of soils with peat 10–40cm deep
- Soils with peaty pockets: areas of mostly non-peat soils, supporting smaller pockets of deep peat (such as flushes or exposures of buried peat) too small to map at a national scale

Peat [substance] – highly organic sediment, composed primarily of partially decomposed plant remains (= subfossils), and preserved through waterlogging

Public value – the value derived by society and people

Proxy indicators – something that is counted/measured in place of a variable of interest, because it responds to that variable in a known way; and that variable can no longer be measured because it happened in the past. A proxy provides an indirect measure of past climates or environments (e.g. moisture levels, temperature, water quality, air pollution etc)

Radiocarbon dating – a radiometric dating technique that uses the radioactive isotope of carbon (¹⁴C)

Raised bog – a peatland which receives its water input solely from rainfall ('ombrotrophic')

Reprofiling – changing the angle of eroded peat to a maximum of either 45 degrees or 33 degrees (depending on follow-up techniques) by the removal (and redeposition) of material

SAC – Special Area of Conservation

SCPs – Spheroidal Carbonaceous Particles; spherical microscopic fly-ash particles, produced from fossil fuel burning

Significance – "The value of a heritage asset to this and future generations because of its heritage interest. That interest may be archaeological, architectural, artistic or historic. Significance derives not only from a heritage asset's physical presence, but also from its setting" (**NPPF** 2019, 64)

SSSI – Site of Special Scientific Interest

Tephra – sherds of volcanic ash emitted during volcanic eruptions. The chemical composition, together with physical characteristics, of tephra layers can be matched to specific eruptions

Testate amoebae – single-celled organisms that indicate moisture levels. They can be used as a proxy to reconstruct past wetness

UNESCO – United Nations Educational, Scientific and Cultural Organization

Value – "An aspect of worth or importance, here attached by people to qualities of places" (English Heritage 2008, 72)

Wasted peat – "a technical term for deep peat that has been substantially degraded following years of drainage and cultivation so that the peat is now more dominated by underlying mineral material" (Natural England 2010, 7)

9. Where to get advice

Local authority curator/archaeologist https://www.algao.org.uk/localgov/hersmr

Local Historic Environment Record (HER) officer https://www.heritagegateway.org.uk/gateway/chr/default.aspx

Historic England's science advisors http://historicengland.org.uk/scienceadvice

Natural England's historic environment advisors https://www.gov.uk/government/publications/natural-englandoffice-locations

10. Acknowledgements

Many thanks to those who have provided input, comments and advice to develop this guidance, in particular: Anna Badcock, Alex Bayliss, Gill Campbell, Rosie Everett, Will Fletcher, Hannah Fluck, Ralph Fyfe, Martin Gillard, Tom Hill, Peter Marshall, Linda Smith, Sue Stallibrass, and the Historic England Science Advisors. Thanks to Sharon Soutar and John Vallender (Historic England) for illustrations and layout design.

Thanks to staff at the Historic England Archive, and to Lee Bray, Jonathan Last, Stephen Macauley, Phil Newman and Nicky Milner for providing photographs.

And finally, many thanks to Louise Brown, Kat Hopwood-Lewis, Naomi Oakley, Mark Phillips and Elaine Willett for helping achieve such a positive outcome from Natural England and Historic England's cross-organisation and cross-sectorial collaborations.

Contact Historic England

East of England Brooklands 24 Brooklands Avenue Cambridge CB2 8BU

Tel: 01223 582749 Email: eastofengland@ HistoricEngland.org.uk

Fort Cumberland Fort Cumberland Road Eastney Portsmouth PO4 9LD

Tel: 023 9285 6704 Email: fort.cumberland@ HistoricEngland.org.uk

London and South East 4th Floor Cannon Bridge House 25 Dowgate Hill London EC4R 2YA

Tel: 020 7973 3700 Email: londonseast@HistoricEngland.org.uk

Midlands The Foundry 82 Granville Street, Birmingham B1 2LH,

Tel: 0121 625 6888 Email: midlands@HistoricEngland.org.uk North East and Yorkshire Bessie Surtees House 41-44 Sandhill Newcastle Upon Tyne NE1 3JF

Tel: 0191 403 1635 Email: northeast@HistoricEngland.org.uk

37 Tanner Row York YO1 6WP

Tel: 01904 601948 Email: yorkshire@HistoricEngland.org.uk

North West 3rd Floor, Canada House 3 Chepstow Street Manchester M1 5FW

Tel: 0161 242 1416 Email: northwest@HistoricEngland.org.uk

South West Fermentation North (1st Floor) Finzels Reach Hawkins Lane Bristol BS1 6JQ

Tel: 0117 975 1308 Email: southwest@HistoricEngland.org.uk

Swindon The Engine House Fire Fly Avenue Swindon SN2 2EH

Tel: 01793 445050 Email: swindon@HistoricEngland.org.uk



We are the public body that helps people care for, enjoy and celebrate England's spectacular historic environment.

Please contact guidance@HistoricEngland.org.uk with any questions about this document.

HistoricEngland.org.uk

If you would like this document in a different format, please contact our customer services department on:

Tel: 0370 333 0607 Email: customers@HistoricEngland.org.uk

All information and weblinks accurate at the time of publication.

Please consider the environment before printing this document