RESEARCH REPORT SERIES no. 14-2013

GODWICK GREAT BARN, GODWICK, TITTLESHALL, NORFOLK TREE-RING DATING OF TIMBERS

SCIENTIFIC DATING REPORT

Alison Arnold and Robert Howard





INTERVENTION AND ANALYSIS

This report has been prepared for use on the internet and the images within it have been down-sampled to optimise downloading and printing speeds.

Please note that as a result of this down-sampling the images are not of the highest quality and some of the fine detail may be lost. Any person wishing to obtain a high resolution copy of this report should refer to the ordering information on the following page. Research Report Series 14-2013

GODWICK GREAT BARN, GODWICK, TITTLESHALL, NORFOLK

TREE-RING DATING OF TIMBERS

Alison Arnold and Robert Howard

NGR: TF 90414 22088

© English Heritage

ISSN 2046-9799 (Print) ISSN 2046-9802 (Online)

The Research Report Series incorporates reports by the expert teams within the Investigation & Analysis Division of the Heritage Protection Department of English Heritage, alongside contributions from other parts of the organisation. It replaces the former Centre for Archaeology Reports Series, the Archaeological Investigation Report Series, the Architectural Investigation Report Series, and the Research Department Report Series.

Many of the Research Reports are of an interim nature and serve to make available the results of specialist investigations in advance of full publication. They are not usually subject to external refereeing, and their conclusions may sometimes have to be modified in the light of information not available at the time of the investigation. Where no final project report is available, readers must consult the author before citing these reports in any publication. Opinions expressed in Research Reports are those of the author(s) and are not necessarily those of English Heritage.

Requests for further hard copies, after the initial print run, can be made by emailing: Res.reports@english-heritage.org.uk or by writing to: English Heritage, Fort Cumberland, Fort Cumberland Road, Eastney, Portsmouth PO4 9LD Please note that a charge will be made to cover printing and postage.

SUMMARY

Samples were taken from the timbers of the roof and floor frames in this building, resulting in the construction and dating of two site sequences and the individual dating of one other sample. Site sequence GDWKSQ01 contains 21 samples and spans the period AD 1406–1597. The second site sequence, GDWKSQ01, contains five samples and spans the period AD 1624–1701. Sample GDW-K20 was found to span the period AD 1582–1683.

The analysis indicates that the extant floor-framing at the northern end of the barn is constructed with timber felled in AD 1597. The majority of the timbers in the roof were felled in AD 1588–1613, though some common rafters were felled in AD 1712–37 and a purlin was potentially felled at the same time or very slightly earlier in AD 1695–1720. These results suggest the floor-frames and primary construction of the roof are contemporary, with both likely to date to, or shortly after AD 1597, with repairs to the roof undertaken in the first decades of the eighteenth century.

CONTRIBUTORS

Alison Arnold and Robert Howard

ACKNOWLEDGEMENTS

The Laboratory would like to thank Mr Garner, owner of the barn, for allowing sampling to be undertaken and Jon Murray of Archaeological Solutions and Stephen Doughty for arranging access. Thanks are also given to Peter Marshall and Cathy Tyers from the English Heritage Scientific Dating Team for their advice and assistance throughout the study and production of this report. Figures used to illustrate the report were provided by Wood and Stephen Design Management.

ARCHIVE LOCATION

Norfolk Historic Environment Service Union House Gressenhall Dereham Norfolk NR20 4DR

DATE OF INVESTIGATION 2012

CONTACT DETAILS

Alison Arnold and Robert Howard Nottingham Tree-ring Dating Laboratory 20 Hillcrest Grove Sherwood Nottingham NG5 1FT roberthoward@tree-ringdating.co.uk alisonarnold@tree-ringdating.co.uk

CONTENTS

Introdu	uction	1
Samplir	רק	1
Analysi	s and Results	1
Interpr	etation	2
Roof.		2
Floor	frames	3
Discus	sion	3
Bibliog	raphy	4
Tables		6
Figures	5	10
Data o	f Measured Samples	24
Append	dix: Tree-Ring Dating	
The P	rinciples of Tree-Ring Dating	30
The P	ractice of Tree-Ring Dating at the Nottingham Tree-Ring Dating Laboratory	30
1.	Inspecting the Building and Sampling the Timbers	30
2.	Measuring Ring Widths	35
3.	Cross-Matching and Dating the Samples	35
4.	Estimating the Felling Date	36
5.	Estimating the Date of Construction	37
6.	Master Chronological Sequences.	
7.	Ring-Width Indices.	
Refer	ences	42

INTRODUCTION

This grade II* listed barn, is located in the deserted medieval village of Godwick, just south of Fakenham, between the villages of Tittleshall and Whissonsett (Figs 1–3). It was associated with the now demolished Godwick Hall which was built in AD 1586. The barn is important due to both its great size and distinctive roof structure. The roof consists of 11 trusses of alternating hammer-beam and queen-strut style (Figs 4 and 5) apart from at the northern end which has as a surviving second floor and attic and evidence of partitioning within the body of the building (Fig 6). This roof is similar to that of the barn at Paston which dates to AD 1581 (Tyers 1999) and at Waxham, dated to AD 1583/4 (Moir 2004). It has been suggested that the construction of these three barns was the result of rivalry amongst the high-status families in the area at this time.

SAMPLING

Sampling was requested by Will Fletcher, English Heritage Inspector of Ancient Monuments, to provide a precise date for this barn that lies within a Scheduled Ancient Monument and to determine its chronological relationship to the barns at Paston and Waxham.

A total of 38 timbers was sampled by coring. Each sample was given the code GDW-K (for Godwick) and numbered 01–38. Twenty-seven of these were from the roof (GDW-K01–27) and 11 from the floor-frames at the northern end of the barn (GDW-K28–38). The location of samples was noted at the time of sampling and has been marked on Figures 7–18. Further details relating to the samples can be found in Table 1. The timber used within the floor frames was of mixed character, a large number being wide ringed and unsuitable for sampling which is why the number of floor samples taken is lower than would perhaps be recommended. Additionally, none of the timbers used within the partitions were thought to be suitable for tree-ring dating, as they clearly contained too few rings for secure dating and so were not sampled.

ANALYSIS AND RESULTS

Ten of the samples, five from the roof and five from the floor frames, were found to have too few rings for secure dating and so were discarded prior to analysis. The remaining 28 samples were prepared by sanding and polishing and their growth-ring widths measured; the data of these measurements are given at the end of the report. These samples were then compared with each other by the Litton/Zainodin grouping procedure (see Appendix), resulting in 26 samples matching to form two groups.

Firstly, 21 samples matched each other and were combined at the relevant offset positions to form GDWKSQ01, a site sequence of 192 rings (Fig 19). This site sequence was compared against a series of relevant reference chronologies where it was found to

match consistently and securely at a first-ring date of AD 1406 and a last-measured ring date of AD 1597. The evidence for this dating is given by the *t*-values in Table 2.

Secondly, five samples grouped and were combined to form GDWKSQ02, a site sequence of 78 rings (Fig 20). When compared against the reference chronologies it was found to span the period AD 1624–1701. The evidence for this dating is given by the *t*-values in Table 3.

Attempts were then made to date the remaining two ungrouped samples by comparing them individually against the reference chronologies resulting in sample GDW-K20 being found to span the period AD 1582–1683. The evidence for this dating is given by the *t*-values in Table 4. The remaining sample, GDW-K19, could not be matched and remains undated.

INTERPRETATION

Tree-ring analysis has resulted in the successful dating of 27 of the samples: 21 from the roof and six from the floor-frames (Fig 21). All felling date ranges have been calculated using the estimate that 95% of mature oak trees in this region have between 15 and 40 sapwood rings.

Roof

There are at least two and potentially three separate fellings represented within the dated timbers from the roof.

Eight samples dated within site sequence GDWKSQ01 have broadly contemporary heartwood/sapwood boundary ring dates, suggestive of a single felling (Figs 19 and 21; Table 1). The average of these is AD 1573, allowing an estimated felling date range to be calculated for the eight timbers represented of AD 1588–1613. The other seven roof samples dated within this site sequence all have last-measured heartwood ring dates in the sixteenth century which, combined with the overall level of similarity found between these ring series, make it likely that they were also felled in AD 1588–1613.

The heartwood/sapwood boundary ring date of four of the samples within site sequence GDWKSQ02 are again suggestive of a single felling (Figs 20 and 21; Table 1). The average of this is AD 1697, allowing an estimated felling date range to be calculated for the four timbers represented of AD 1712–37. These four timbers are all common rafters and it seems likely that the other dated common rafter within site sequence GDWKSQ02, with a last-measured ring date of AD 1689 was also likely to have been felled in AD 1712–37.

The individually dated sample GDW-K20, taken from a purlin, has a heartwood/sapwood boundary ring date of AD 1680 (Fig 21; Table 1), allowing an estimated felling date range to be calculated for the timber represented of AD 1695–1720.

Floor frames

Six of these samples, representing two main beams from the first floor and six joists from the second floor, have been dated within site sequence GDWKSQ01. Two of these timbers, a first-floor main beam and a second-floor joist, have complete sapwood and the last-measured ring date of AD 1597, the felling date of the two timbers represented. Three of the remaining dated timbers have heartwood/sapwood boundary dates which are similar and imply that these are also likely to have been felled in AD 1597. With a last measured heartwood ring date of AD 1570 it appears likely that the last of the dated samples from the floor-frames is likely to be contemporary and hence probably felled in AD 1597.

DISCUSSION

Prior to tree-ring analysis being undertaken on the timbers of the roof and floors at this barn it was thought likely to date to sometime shortly after AD 1586, the date of construction of Godwick Hall. The similarity in design between this roof and those of the great barns at Paston and Waxham had been highlighted but it was unclear where this barn fitted chronologically.

It is now known that the majority of the dated timbers from the roof were felled in AD 1588–1613, whilst the floor frames contain timber felled in AD 1597. These results suggest that the roof and floor frames are contemporary, with the construction of both likely to have occurred in or soon after felling of the timbers in AD 1597. This thus places Godwick Great barn slightly later than the barns at Paston (AD 1581) and Waxham (AD 1583/4).

In addition a number of common rafters and a purlin clearly represent later repairs to the roof just over a century after the initial construction. The common rafters were felled in the period AD 1712–37, whilst the purlin has an estimated felling date of AD 1695–1720. Clearly it is possible that these six timbers were all part of a single felling episode in the first decades of the eighteenth century.

BIBLIOGRAPHY

Arnold, A J, Howard, R E, Laxton, R R, and Litton, C D, 2002 *The Urban Development of Newark-on-Trent: A Dendrochronological Approach*, Centre for Archaeol Rep, **95/2002**

Arnold, A J, and Howard, R E, 2003 Tree-ring analysis of samples from Cobham Hall, Cobham, Kent, unpubl composite site sequence *COBBCDHFSQ01*, NTRDL

Arnold, A J, Howard, R E, and Litton, C D, 2003a *Tree-ring analysis of timbers from Manor House, West Street, Alford, Lincolnshire,* Centre for Archaeol Rep, **55/2003**

Arnold, A J, Howard, R E, and Litton, C D, 2003b *Tree-ring analysis of timbers from Hulme Hall, Allostock, near Northwich*, Centre for Archaeol Rep, **84/2003**

Arnold, A J, Howard, R E, Laxton, R R, Litton, C D, and Simpson, W G, 2004 *A Dendrochronological study of the Monastic Buildings of Ely*, Centre for Archaeol Rep, **74/2004**

Arnold, A J, and Howard, R E, 2006 Tree-ring analysis of timbers from Newnham Hall Farm, Newnham Murren, near Wallingford, Oxfordshire, unpubl computer files *CMGASQ01/CMGBSQ02*, NTRDL

Arnold, A J, and Howard, R E, 2007 Nottingham Tree-ring Dating Laboratory: results, *Vernacular Architect*, **38**, 106–14

Arnold, A J, Howard, R E, and Litton, C D, 2008 *Shenton Hall, Shenton, Leicestershire, treering analysis of timbers from the dovecote,* English Heritage Res Dep Rep Ser, **22/2008**

Arnold, A J, and Howard, R E, 2009 Tree-ring analysis of timbers from Castle Street, Melbourne, Derbyshire, unpublisite sequence *MLBDSQ01*, NTRDL

Arnold, A J, and Howard, R E, 2010a *St Firmin Church, Thurlby, Lincolnshire; Tree-Ring Analysis of Timbers of the Bellframe and Tower*, English Heritage Res Dep Rep Ser, **72/2010**

Arnold, A J, and Howard, R E, 2010b *All Saints' Church, North Scarle, Lincolnshire: Tree-Ring Analysis of Timbers of the Bellframe and Belfry Floor*, English Heritage Res Dep Rep Ser, **44/2010**

Howard, R E, and Litton, C D, 1997 *Tree-ring analysis of timbers from Astley Castle, Warwickshire*, Anc Mon Lab Rep, **83/1997**

Howard, R E, Laxton, R R, and Litton, C D, 2005 *Tree-ring analysis of timbers from the Riding School, Bolsover Castle, Bolsover, Derbyshire*, Centre for Archaeol Rep, **40/2005**

Laxton, R R, Litton, C D, Simpson, W G, and Whitley, J P, 1982 Tree-ring Dates from some East Midlands Buildings: Table 1, no 7, *Trans of the Thoroton Society of Nottinghamshire*, **86**, 76–7

Laxton, R R, and Litton, C D, 1988 *An East Midlands master tree-ring chronology and its use for dating vernacular buildings*, Univ of Nottingham Dept of Classical and Archaeol Studies, Monograph Series, **III**

Miles, D, 2007 H M Tower of London (Tol99 & Tol100), *London Borough of Tower Hamlets : The Tree-Ring Dating of the White Tower*, Res Dep Rep, **35/2007**

Miles D, and Worthington, M, 1997 Tree-ring Dates, Vernacular Architect, 28, 159–81

Miles D, and Worthington, M, 2006 *Tree-ring dating of the Fellow's Quadrangle, Merton College, Oxford, Oxfordshire*, Res Dep Rep, **80/2006**

Moir, A, 2004 *Tree-ring Analysis of Timbers from Waxham Great Barn, Sea Palling, Norfolk,* Centre for Archaeol Rep, **67/2004**

Tyers, I, 1997 *Tree-ring Analysis of Timbers from Sinai Park, Staffordshire*, Anc Mon Lab Rep, **80/97**

Tyers, I, 1999 *Tree-ring Analysis of Timbers from Paston Great Barn, Norfolk*, Anc Mon Lab Rep, **54/1999**

TABLES

Table 1: Details of tree-ring samples from Godwick Great Barn, Tittleshall, Norfolk

Sample	Sample location	Total rings*	Sapwood rings**	First measured ring	Last heartwood ring	Last measured ring	
number				date (AD)	date (AD)	date (AD)	
Roof							
GDW-K01	Collar, truss 1	NM					
GDW-K02	East middle purlin, bay 1	NM					
GDW-K03	West middle purlin, bay 1	119		1434		1552	
GDW-K04	East strut, truss 2	NM					
GDW-K05	West strut, truss 2	73	01	1509	1580	1581	
GDW-K06	East principal rafter, truss 3	NM					
GDW-K07	East corbel, truss 4	54	h/s	1517	1570	1570	
GDW-K08	East principal rafter, truss 5	50		1518		1567	
GDW-K09	West common rafter 2, bay 5	74	h/s	1627	1700	1700	
GDW-K10	East common rafter 3, bay 5	63	h/s	1630	1692	1692	
GDW-K11	East common rafter 4, bay 5	58		1632		1689	
GDW-K12	West common rafter 5, bay 5	71	02	1630	1698	1700	
GDW-K13	West hammer beam, truss 6	55	01	1528	1581	1582	
GDW-K14	East hammer beam, truss 6	NM					
GDW-K15	West corbel, truss 6	55	h/s	1514	1568	1568	
GDW-K16	East common rafter 2, bay 6	78	04	1624	1697	1701	
GDW-K17	East corbel, truss 7	53	h/s	1517	1569	1569	
GDW-K18	East hammer beam, truss 8	79		1490		1568	
GDW-K19	West hammer beam, truss 8	57	h/s				
GDW-K20	East lower purlin, bay 8	102	03	1582	1680	1683	
GDW-K21	East common rafter 2, bay 8	54		1500		1553	
GDW-K22	East common rafter 4, bay 8	96		1429		1524	
GDW-K23	East principal rafter, truss 9	101		1422		1522	
GDW-K24	West principal rafter, truss 9	169	h/s	1406	1574	1574	
GDW-K25	West corbel, truss 9	57	02	1514	1568	1570	
GDW-K26	East hammer beam, truss 10	92		1481		1572	

6

GDW-K27	West hammer beam, truss 10	82	h/s	1494	1575	1575	
Floor		·	•		•		
GDW-K28	1st floor, main beam, truss 2	68	17C	1530	1580	1597	
GDW-K29	1st floor, main beam, truss 3	50	h/s	1531	1580	1580	
GDW-K30	2nd floor, main beam, truss 2	NM					
GDW-K31	2nd floor, joist 11, bay 1	60		1511		1570	
GDW-K32	2nd floor, joist 5, bay 1	NM					
GDW-K33	2nd floor, joist 9, bay 1	67	17C	1531	1580	1597	
GDW-K34	2nd floor, joist 13, bay 1	58	h/s	1523	1580	1580	
GDW-K35	2nd floor, joist 16, bay 1	NM					
GDW-K36	2nd floor, joist 19, bay 1	56	h/s	1514	1569	1569	
GDW-K37	2nd floor, joist 14, bay 2	NM					
GDW-K38	2nd floor, joist 15, bay 2	NM					

*NM = not measured

**h/s = the heartwood/sapwood boundary is the last measured ring on the sample C = complete sapwood retained on sample, last-measured ring is the felling date

Table 2: Results of the cross-matching of site sequence GDWKSQ01 and the reference chronologies when the first-ring date is AD 1406 and the last measured ring date is AD 1597

Reference chronology	<i>t</i> -value	Span of chronology	Reference
Newnham Hall Farm House, Newnham Murren, near Wallingford, Oxfordshire	9.7	AD 1412-1614	Arnold and Howard 2006 unpubl
Newnham Hall Farm Cottage, Newnham Murren, near Wallingford, Oxfordshire	8.4	AD 1414–1551	Arnold and Howard 2006 unpubl
Cobham Hall, Cobham, Kent	7.6	AD 1317–1663	Arnold and Howard 2003 unpubl
The White Tower, Tower of London, London	7.5	AD 1463-1612	Miles 2007
Powcher's Hall (east roof), Ely Cathedral Precinct, Ely, Cambridgeshire	7.3	AD 1457-1609	Arnold et al 2004
Manor House, Alford, Lincolnshire	7.1	AD 1500-1668	Arnold et al 2003a
Astley Castle, Warwickshire	7.0	AD 1495–1627	Howard and Litton 1997

Table 3: Results of the cross-matching of site sequence GDWKSQ02 and the reference chronologies when the first-ring date is AD 1624 and the last measured ring date is AD 1701

Reference chronology	<i>t</i> -value	Span of chronology	Reference
East Midlands	6.7	AD 882–1981	Laxton and Litton 1988
Castle House, Melbourne, Derbyshire	6.4	AD 1583–1720	Arnold and Howard 2009 unpubl
Shenton Hall Dovecote, Shenton, Leicestershire	6.2	AD 1606–1719	Arnold et al 2008
St Firmin Church, Thurlby, Lincolnshire	5.8	AD 1599–1792	Arnold and Howard, 2010a
All Saints' Church, North Scarle, Lincolnshire	5,8	AD 1602–1716	Arnold and Howard 2010b
6/12 Chain Lane, Newark, Nottinghamshire	5.6	AD 1608–1684	Arnold et al 2002
Sinai Park, Burton on Trent, Staffordshire	5.6	AD 1227–1750	Tyers 1997

Table 4: Results of the cross-matching of sample GDW-K20 and the reference chronologies when the first-ring date is AD 1582 and the last measured ring date is AD 1683

Reference chronology	<i>t</i> -value	Span of chronology	Reference
-			
East Midlands	6.3	AD 882–1981	Laxton and Litton 1988
Blidworth, Nottinghamshire	5.9	AD 1621–1713	Laxton <i>et al</i> 1982
Bolsover Castle (Riding School), Bolsover, Derbyshire	5.2	AD 1494–1744	Howard et al 2005
Fellows Quad, Mertons College, Oxfordshire	5.2	AD 1442–1608	Miles and Worthington 2006
Hulme Hall, Allostock, Near Northwich,	5.1	AD 1574–1689	Arnold et al 2003b
Hempshill Hall, Nottinghamshire	5.1	AD 1566–1702	Arnold and Howard 2007
Stokesay Castle, Shropshire	4.9	AD 1463–1662	Miles and Worthington 1997

FIGURES




Figure 1: Map to show the general location of Godwick Great Barn, arrowed. © *Crown Copyright and database right 2013. All rights reserved. Ordnance Survey Licence number 100024900*



Figure 2: Map to show the location of Godwick Great Barn, hatched. © *Crown Copyright and database right 2013. All rights reserved. Ordnance Survey Licence number 100024900*



Figure 3: Godwick Great Barn roof (photograph taken from the south-west) Photograph Robert Howard






Figure 4: Ground-floor plan; showing truss locations (Wood and Stephen Design Management)




Figure 5: Godwick Great Barn partitioning at truss 5 in the foreground (photograph taken from the south-west) Photograph Robert Howard




Figure 6: Truss 1, showing the location of samples GDW-K01–3 (Wood and Stephen Design Management)




Figure 7: Truss 2, showing the location of samples GDW-K04, GDW-K05, and GDW-K28 (Wood and Stephen Design Management)




Figure 8: Truss 3, showing the location of samples GDW-K06 and GDW-K29 (Wood and Stephen Design Management)




Figure 9: Truss 4, showing the location of sample GDW-K07 (Wood and Stephen Design Management)




Figure 10: Truss 5, showing the location of sample GDW-K08 (Wood and Stephen Design Management)




Figure 11: Truss 6, showing the location of samples GDW-K13–15 (Wood and Stephen Design Management)




Figure 12: Truss 7, showing the location of sample GDW-K17 (Wood and Stephen Design Management)




Figure 13: Truss 8, showing the location of samples GDW-K18-20 (Wood and Stephen Design Management)




Figure 14: Truss 9, showing the location of samples GDW-K23-5 (Wood and Stephen Design Management)




Figure 15: Truss 10, showing the location of samples GDW-K26 and GDW-K27 (Wood and Stephen Design Management)




Figure 16: Plan of rafters, showing the location of samples GDW-K09-12, GDW-K16, and GDW-K21-2 (after Wood and Stephen Design Management)





Figure 17: Second-floor plan, showing the location of samples GDW-K30-38 (after Wood and Stephen Design Management)



 $^{\odot}$

Figure 18: Bar diagram of samples in site sequence GDWKSQ01




Figure 19: Bar diagram of samples in site sequence GDWKSQ02

© ENGLISH HERITAGE

22

14 - 2013




Figure 20: Bar diagram of all dated samples, sorted by heartwood/sapwood boundary and area

DATA OF MEASURED SAMPLES

Measurements in 0.01mm units

352 260 280 217 366 234 180 203 375 418 240 97 154 216 341 235 247 296 412 321 341 268 319 242 269 119 130 144 224 232 200 143 118 68 78 112 105 104 147 137 107 109 85 157 201 144 132 166 154 218 162 152 129 126 197 152 86 64 118 101 105 152 175 140 162 141 110

GDW-K34A 58

174 210 147 157 228 193 238 137 232 200 176 164 212 164 194 204 270 367 182 80 103 117 166 156 228 253 237 191 151 167 205 166 144 90 65 91 84 125 118 134 115 83 113 126 100 152 90 123 87 65 91 113 138 97 96 107 132 148 GDW-K34B 58

157 211 152 154 192 197 253 149 205 188 164 212 199 165 208 219 267 363 188 82 99 118 167 159 226 252 252 182 167 172 200 166 145 91 61 86 94 110 109 139 112 101 106 122 111 151 92 124 86 71 91 120 145 99 101 95 126 140 GDW-K36A 56

304 244 315 350 468 402 253 210 175 118 173 213 175 122 102 80 67 190 432 557 560 426 382 457 431 602 505 229 105 202 265 301 379 338 361 459 374 299 249 382 256 242 114 122 157 238 251 396 283 284 157 136 122 104 137 180 GDW-K36B 56

300 251 308 336 467 416 241 211 178 117 189 205 177 128 96 78 63 196 452 512 522 424 392 448 444 592 506 228 117 188 276 299 364 343 355 469 358 316 252 387 249 238 119 126 157 232 238 410 281 276 169 128 112 110 144 196

APPENDIX: TREE-RING DATING

The Principles of Tree-Ring Dating

Tree-ring dating, or dendrochronology as it is known, is discussed in some detail in the Laboratory's Monograph, An East Midlands Master Tree-Ring Chronology and its uses for dating Vernacular Building (Laxton and Litton 1988) and Dendrochronology: Guidelines on Producing and Interpreting Dendrochronological Dates (English Heritage 1998). Here we will give the bare outlines. Each year an oak tree grows an extra ring on the outside of its trunk and all its branches just inside its bark. The width of this annual ring depends largely on the weather during the growing season, about April to October, and possibly also on the weather during the previous year. Good growing seasons give rise to relatively wide rings, poor ones to very narrow rings and average ones to relatively average ring widths. Since the climate is so variable from year to year, almost randomlike, the widths of these rings will also appear random-like in sequence, reflecting the seasons. This is illustrated in Figure A1 where, for example, the widest rings appear at irregular intervals. This is the key to dating by tree rings, or rather, by their widths. Records of the average ring widths for oaks, one for each year for the last 1000 years or more, are available for different areas. These are called master chronologies. Because of the random-like nature of these sequences of widths, there is usually only one position at which a sequence of ring widths from a sample of oak timber with at least 70 rings will match a master. This will date the timber and, in particular, the last ring.

If the bark is still on the sample, as in Figure A1, then the date of the last ring will be the date of felling of the oak from which it was cut. There is much evidence that in medieval times oaks cut down for building purposes were used almost immediately, usually within the year or so (Rackham 1976). Hence if bark is present on several main timbers in a building, none of which appear reused or are later insertions, and if they all have the same date for their last ring, then we can be quite confident that this is the date of construction or soon after. If there is no bark on the sample, then we have to make an estimate of the felling date; how this is done is explained below.

The Practice of Tree-Ring Dating at the Nottingham Tree-Ring Dating Laboratory

1. Inspecting the Building and Sampling the Timbers. Together with a building historian the timbers in a building are inspected to try to ensure that those sampled are not reused or later insertions. Sampling is almost always done by coring into the timber, which has the great advantage that we can sample in situ timbers and those judged best to give the date of construction, or phase of construction if there is more than one in the building. The timbers to be sampled are also inspected to see how many rings they have. We normally look for timbers with at least 70 rings, and preferably more. With fewer rings than this, 50 for example, sequences of widths become difficult to match to a unique


position within a master sequence of ring widths and so are difficult to date (Litton and Zainodin 1991). The cross-section of the rafter shown in Figure A2 has about 120 rings; about 20 of which are sapwood rings – the lighter rings on the outside. Similarly the core has just over 100 rings with a few sapwood rings.

To ensure that we are getting the date of the building as a whole, or the whole of a phase of construction if there is more than one, about 8–10 samples per phase are usually taken. Sometimes we take many more, especially if the construction is complicated. One reason for taking so many samples is that, in general, some will fail to give a date. There may be many reasons why a particular sequence of ring widths from a sample of timber fails to give a date even though others from the same building do. For example, a particular tree may have grown in an odd ecological niche, so odd indeed that the widths of its rings were determined by factors other than the local climate! In such circumstances it will be impossible to date a timber from this tree using the master sequence whose widths, we can assume, were predominantly determined by the local climate at the time.

Sampling is done by coring into the timber with a hollow corer attached to an electric drill and usually from its outer rings inwards towards where the centre of the tree, the pith, is judged to be. An illustration of a core is shown in Figure A2; it is about 150mm long and 10mm diameter. Great care has to be taken to ensure that as few as possible of the outer rings are lost in coring. This can be difficult as these outer rings are often very soft (see below on sapwood). Each sample is given a code which identifies uniquely which timber it comes from, which building it is from and where the building is located. For example, CRO-A06 is the sixth core taken from the first building (A) sampled by the Laboratory in Cropwell Bishop. Where it came from in that building will be shown in the sampling records and drawings. No structural damage is done to any timbers by coring, nor does it weaken them.

During the initial inspection of the building and its timbers the dendrochronologist may come to the conclusion that, as far as can be judged, none of the timbers have sufficient rings in them for dating purposes and may advise against sampling to save further unwarranted expense.

All sampling by the Laboratory is undertaken according to current Health and Safety Standards. The Laboratory's dendrochronologists are insured.



on the outside just inside the bark. The year of each ring can be determined by counting back from the outside ring, which grew in 1976



Figure A2: Cross-section of a rafter, showing sapwood rings in the left-hand corner, the arrow points to the heartwood/sapwood boundary (H/S); and a core with sapwood; again the arrow is pointing to the H/S. The core is about the size of a pencil



Figure A3: Measuring ring widths under a microscope. The microscope is fixed while the sample is on a moving platform. The total sequence of widths is measured twice to ensure that an error has not been made. This type of apparatus is needed to process a large number of samples on a regular basis



Figure A4: Three cores from timbers in a building. They come from trees growing at the same time. Notice that, although the sequences of widths look similar, they are not identical. This is typical

2. Measuring Ring Widths. Each core is sanded down with a belt sander using medium-grit paper and then finished by hand with flourgrade-grit paper. The rings are then clearly visible and differentiated from each other with a result very much like that shown in Figure A2. The core is then mounted on a movable table below a microscope and the ring-widths measured individually from the innermost ring to the outermost. The widths are automatically recorded in a computer file as they are measured (see Fig A3).

3. Cross-Matching and Dating the Samples. Because of the factors besides the local climate which may determine the annual widths of a tree's rings, no two sequences of ring widths from different oaks growing at the same time are exactly alike (Fig A4). Indeed, the sequences may not be exactly alike even when the trees are growing near to each other. Consequently, in the Laboratory we do not attempt to match two sequences of ring widths by eye, or graphically, or by any other subjective method. Instead, it is done objectively (ie statistically) on a computer by a process called cross-matching. The output from the computer tells us the extent of correlation between two sample sequences of widths or, if we are dating, between a sample sequence of widths and the master, at each relative position of one to the other (offsets). The extent of the correlation at an offset is determined by the t-value (defined in almost any introductory book on statistics). That offset with the maximum t-value among the t-values at all the offsets will be the best candidate for dating one sequence relative to the other. If one of these is a master chronology, then this will date the other. Experiments carried out in the past with sequences from oaks of known date suggest that a t-value of at least 4.5, and preferably at least 5.0, is usually adequate for the dating to be accepted with reasonable confidence (Laxton and Litton 1988; Laxton et al 1988; Howard et al 1984–1995).

This is illustrated in Figure A5 with timbers from one of the roofs of Lincoln Cathedral. Here four sequences of ring widths, LIN-CO4, 05, 08, and 45, have been cross-matched with each other. The ring widths themselves have been omitted in the bar diagram, as is usual, but the offsets at which they best cross-match each other are shown; eg the sequence of ring widths of CO8 matches the sequence of ring widths of C45 best when it is at a position starting 20 rings after the first ring of C45, and similarly for the others. The actual t-values between the four at these offsets of best correlations are in the matrix. Thus at the offset of +20 rings, the t-value between C45 and C08 is 5.6 and is the maximum found between these two among all the positions of one sequence relative to the other.

It is standard practice in our Laboratory first to cross-match as many as possible of the ring-width sequences of the samples in a building and then to form an average from them. This average is called a site sequence of the building being dated and is illustrated in Figure A5. The fifth bar at the bottom is a site sequence for a roof at Lincoln Cathedral and is constructed from the matching sequences of the four timbers. The site sequence width for each year is the average of the widths in each of the sample sequences which has a width for that year. Thus in Fig A5 if the widths shown are 0.8mm for C45, 0.2mm for C08, 0.7mm for C05, and 0.3mm for C04, then the corresponding width of the site

sequence is the average of these, 0.55mm. The actual sequence of widths of this site sequence is stored on the computer. The reason for creating site sequences is that it is usually easier to date an average sequence of ring widths with a master sequence than it is to date the individual component sample sequences separately.

The straightforward method of cross-matching several sample sequences with each other one at a time is called the 'maximal t-value' method. The actual method of cross-matching a group of sequences of ring-widths used in the Laboratory involves grouping and averaging the ring-width sequences and is called the 'Litton-Zainodin Grouping Procedure'. It is a modification of the straightforward method and was successfully developed and tested in the Laboratory and has been published (Litton and Zainodin 1991; Laxton et al 1988).

4. Estimating the Felling Date. As mentioned above, if the bark is present on a sample, then the date of its last ring is the date of the felling of its tree (or the last full year before felling, if it was felled in the first three months of the following calendar year, before any new growth had started, but this is not too important a consideration in most cases). The actual bark may not be present on a timber in a building, though the dendrochronologist who is sampling can often see from its surface that only the bark is missing. In these cases the date of the last ring is still the date of felling.

Quite often some, though not all, of the original outer rings are missing on a timber. The outer rings on an oak, called sapwood rings, are usually lighter than the inner rings, the heartwood, and so are relatively easy to identify. For example, sapwood can be seen in the corner of the rafter and at the outer end of the core in Figure A2, both indicated by arrows. More importantly for dendrochronology, the sapwood is relatively soft and so liable to insect attack and wear and tear. The builder, therefore, may remove some of the sapwood for precisely these reasons. Nevertheless, if at least some of the sapwood rings are left on a sample, we will know that not too many rings have been lost since felling so that the date of the last ring on the sample is only a few years before the date of the original last ring on the tree, and so to the date of felling.

Various estimates have been made and used for the average number of sapwood rings in mature oak trees (English Heritage 1998). A fairly conservative range is between 15 and 50 and that this holds for 95% of mature oaks. This means, of course, that in a small number of cases there could be fewer than 15 and more than 50 sapwood rings. For example, the core CRO-A06 has only 9 sapwood rings and some have obviously been lost over time – either they were removed originally by the carpenter and/or they rotted away in the building and/or they were lost in the coring. It is not known exactly how many sapwood rings are missing, but using the above range the Laboratory would estimate between a minimum of 6 (=15-9) and a maximum of 41 (=50-9). If the last ring of CRO-A06 has been dated to 1500, say, then the estimated felling-date range for the tree from which it came originally would be between 1506 and 1541. The Laboratory uses this estimate for sapwood in areas of England where it has no prior information. It

also uses it when dealing with samples with very many rings, about 120 to the last heartwood ring. But in other areas of England where the Laboratory has accumulated a number of samples with complete sapwood, that is, no sapwood lost since felling, other estimates in place of the conservative range of 15 to 50 are used. In the East Midlands (Laxton et al 2001) and the east to the south down to Kent (Pearson 1995) where it has sampled extensively in the past, the Laboratory uses the shorter estimate of 15 to 35 sapwood rings in 95% of mature oaks growing in these parts. Since the sample CRO-A06 comes from a house in Cropwell Bishop in the East Midlands, a better estimate of sapwood rings lost since felling is between a minimum of 6 (=15-9) and 26 (=35-9) and the felling would be estimated to have taken place between 1506 and 1526, a shorter period than before. Oak boards quite often come from the Baltic region and in these cases the 95% confidence limits for sapwood are 9 to 36 (Howard et al 1992, 56).

Even more precise estimates of the felling date and range can often be obtained using knowledge of a particular case and information gathered at the time of sampling. For example, at the time of sampling the dendrochronologist may have noted that the timber from which the core of Figure A2 was taken still had complete sapwood but that some of the soft sapwood rings were lost in coring. By measuring into the timber the depth of sapwood lost, say 20mm, a reasonable estimate can be made of the number of sapwood rings lost, say 12 to 15 rings in this case. By adding on 12 to 15 years to the date of the last ring on the sample a good tight estimate for the range of the felling date can be obtained, which is often better than the 15 to 35 years later we would have estimated without this observation. In the example, the felling is now estimated to have taken place between AD 1512 and 1515, which is much more precise than without this extra information.

Even if all the sapwood rings are missing on a sample, but none of the heartwood rings are, then an estimate of the felling-date range is possible by adding on the full compliment of, say, 15 to 35 years to the date of the last heartwood ring (called the heartwood/ sapwood boundary or transition ring and denoted H/S). Fortunately it is often easy for a trained dendrochronologist to identify this boundary on a timber. If a timber does not have its heartwood/sapwood boundary, then only a post quem date for felling is possible.

5. Estimating the Date of Construction. There is a considerable body of evidence collected by dendrochronologists over the years that oak timbers used in buildings were not seasoned in medieval or early modern times (English Heritage 1998; Miles 1997, 50–5). Hence, provided that all the samples in a building have estimated felling-date ranges broadly in agreement with each other, so that they appear to have been felled as a group, then this should give an accurate estimate of the period when the structure was built, or soon after (Laxton et al 2001, fig 8; 34–5, where 'associated groups of fellings' are discussed in detail). However, if there is any evidence of storage before use, or if there is evidence the oak came from abroad (eg Baltic boards), then some allowance has to be made for this.

6. Master Chronological Sequences. Ultimately, to date a sequence of ring widths, or a site sequence, we need a master sequence of dated ring widths with which to crossmatch it, a Master Chronology. To construct such a sequence we have to start with a sequence of widths whose dates are known and this means beginning with a sequence from an oak tree whose date of felling is known. In Figure A6 such a sequence is SHE-T, which came from a tree in Sherwood Forest which was blown down in a recent gale. After this other sequences which cross-match with it are added and gradually the sequence is 'pushed back in time' as far as the age of samples will allow. This process is illustrated in Figure A6. We have a master chronological sequence of widths for Nottinghamshire and East Midlands oak for each year from AD 882 to 1981. It is described in great detail in Laxton and Litton (1988), but the components it contains are shown here in the form of a bar diagram. As can be seen, it is well replicated in that for each year in this period there are several sample sequences having widths for that year. The master is the average of these. This master can now be used to date oak from this area and from the surrounding areas where the climate is very similar to that in the East Midlands. The Laboratory has also constructed a master for Kent (Laxton and Litton 1989). The method the Laboratory uses to construct a master sequence, such as the East Midlands and Kent, is completely objective and uses the Litton-Zainodin grouping procedure (Laxton et al 1988). Other laboratories and individuals have constructed masters for other areas and have made them available. As well as these masters, local (dated) site chronologies can be used to date other buildings from nearby. The Laboratory has hundreds of these site sequences from many parts of England and Wales covering many short periods.

7. **Ring-Width Indices.** Tree-ring dating can be done by cross-matching the ring widths themselves, as described above. However, it is advantageous to modify the widths first. Because different trees grow at different rates and because a young oak grows in a different way from an older oak, irrespective of the climate, the widths are first standardized before any matching between them is attempted. These standard widths are known as ring-width indices and were first used in dendrochronology by Baillie and Pilcher (1973). The exact form they take is explained in this paper and in the appendix of Laxton and Litton (1988) and is illustrated in the graphs in Figure A7. Here ring-widths are plotted vertically, one for each year of growth. In the upper sequence of (a), the generally large early growth after 1810 is very apparent as is the smaller later growth from about 1900 onwards when the tree is maturing. A similar phenomenon can be observed in the lower sequence of (a) starting in 1835. In both the widths are also changing rapidly from year to year. The peaks are the wide rings and the troughs are the narrow rings corresponding to good and poor growing seasons, respectively. The two corresponding sequence of Baillie-Pilcher indices are plotted in (b) where the differences in the immature and mature growths have been removed and only the rapidly changing peaks and troughs remain, that are associated with the common climatic signal. This makes cross-matching easier.

t-value/offset Matrix




Figure A5: Cross-matching of four sequences from a Lincoln Cathedral roof and the formation of a site sequence from them

The bar diagram represents these sequences without the rings themselves. The length of the bar is proportional to the number of rings in the sequence. Here the four sequences are set at relative positions (offsets) to each other at which they have maximum correlation as measured by the t-values. The t-value/offset matrix contains the maximum t-values below the diagonal and the offsets above it. Thus, the maximum t-value between C08 and C45 occurs at the offset of +20 rings and the t-value is then 5.6. The site sequence is composed of the average of the corresponding widths, as illustrated with one width.

14 - 2013




Figure A6: Bar diagram showing the relative positions and dates of the first rings of the component site sequences in the East Midlands Master Dendrochronological Sequence, EM08/87




Figure A7 (a): The raw ring-widths of two samples, THO-A01 and THO-B05, whose felling dates are known

Here the ring widths are plotted vertically, one for each year, so that peaks represent wide rings and troughs narrow ones. Notice the growth-trends in each; on average the earlier rings of the young tree are wider than the later ones of the older tree in both sequences

Figure A7 (b): The Baillie-Pilcher indices of the above widths

The growth trends have been removed completely

References

Baillie, M G L, and Pilcher, J R, 1973 A simple cross-dating program for tree-ring research, *Tree-Ring Bull*, **33**, 7–14

English Heritage, 1998 Dendrochronology: *Guidelines on Producing and Interpreting Dendrochronological Dates*, London

Hillam, J, Morgan, R A, and Tyers, I, 1987 Sapwood estimates and the dating of short ring sequences, *Applications of tree-ring studies*, BAR Int Ser, **3**, 165–85

Howard, R E, Laxton, R R, Litton, C D, and Simpson, W G, 1984–95 Nottingham University Tree-Ring Dating Laboratory results, *Vernacular Architect*, **15–26**

Howard, R E, Laxton, R R, Litton, C D, and Simpson, W G, 1992 List 44 no 17 -Nottingham University Tree-Ring Dating Laboratory: tree-ring dates for buildings in the East Midlands, *Vernacular Architect*, **23**, 51–6

Hughes, M K, Milson, S J, and Legett, P A, 1981 Sapwood estimates in the interpretation of tree-ring dates, *J Archaeol Sci*, **8**, 381–90

Laxon, R R, Litton, C D, and Zainodin, H J, 1988 An objective method for forming a master ring-width sequence, *P A C T*, **22**, 25–35

Laxton, R R, and Litton, C D, 1988 *An East Midlands Master Chronology and its use for dating vernacular buildings*, University of Nottingham, Department of Archaeology Publication, Monograph Series III

Laxton, R R, and Litton, C D, 1989 Construction of a Kent master dendrochronological sequence for oak, AD 1158 to 1540, *Medieval Archaeol*, **33**, 90–8

Laxton, R R, Litton, C D, and Howard, R E, 2001 *Timber: Dendrochronology of Roof Timbers at Lincoln Cathedral*, Engl Heritage Res Trans, **7**

Litton, C D, and Zainodin, H J, 1991 Statistical models of dendrochronology, *J Archaeol Sci*, **18**, 29–40

Miles, D W H, 1997 The interpretation, presentation and use of tree-ring dates, *Vernacular Architect*, **28**, 40–56

Pearson, S, 1995 The Medieval Houses of Kent, an Historical Analysis, London

Rackham, O, 1976 Trees and Woodland in the British Landscape, London



ENGLISH HERITAGE RESEARCH AND THE HISTORIC ENVIRONMENT

English Heritage undertakes and commissions research into the historic environment, and the issues that affect its condition and survival, in order to provide the understanding necessary for informed policy and decision making, for the protection and sustainable management of the resource, and to promote the widest access, appreciation and enjoyment of our heritage. Much of this work is conceived and implemented in the context of the National Heritage Protection Plan. For more information on the NHPP please go to http://www.english-heritage. org.uk/professional/protection/national-heritage-protection-plan/.

The Heritage Protection Department provides English Heritage with this capacity in the fields of building history, archaeology, archaeological science, imaging and visualisation, landscape history, and remote sensing. It brings together four teams with complementary investigative, analytical and technical skills to provide integrated applied research expertise across the range of the historic environment. These are:

- * Intervention and Analysis (including Archaeology Projects, Archives, Environmental Studies, Archaeological Conservation and Technology, and Scientific Dating)
- * Assessment (including Archaeological and Architectural Investigation, the Blue Plaques Team and the Survey of London)
- * Imaging and Visualisation (including Technical Survey, Graphics and Photography)
- * Remote Sensing (including Mapping, Photogrammetry and Geophysics)

The Heritage Protection Department undertakes a wide range of investigative and analytical projects, and provides quality assurance and management support for externally-commissioned research. We aim for innovative work of the highest quality which will set agendas and standards for the historic environment sector. In support of this, and to build capacity and promote best practice in the sector, we also publish guidance and provide advice and training. We support community engagement and build this in to our projects and programmes wherever possible.

We make the results of our work available through the Research Report Series, and through journal publications and monographs. Our newsletter *Research News*, which appears twice a year, aims to keep our partners within and outside English Heritage up-to-date with our projects and activities.

A full list of Research Reports, with abstracts and information on how to obtain copies, may be found on www.english-heritage.org.uk/researchreports

For further information visit www.english-heritage.org.uk

